arXiv:0805.0458v1 [math.CA] 5 May 2008

The Elliptic Hypergeometric Functions Associated to the Configuration Space of Points on an Elliptic Curve I : Twisted Cycles

Ko-Ki Ito

May 5, 2008

Abstract

We consider the Euler type integral associated to the configuration space of points on an elliptic curve, which is an analogue of the hypergeometric function associated to the configuration space of points on a projective line. We calculate the *twisted homology group*, with coefficients in the local system associated to a power function g^{α} of an elliptic function g, and the intersection form. Applying these calculations, we describe the *connection matrices* representing the linear isomorphisms induced from analytic continuations of the functions defined by the integrations of g^{α} over twisted cycles.

Key Words and Phrases: elliptic curve, twisted homology, Euler type integral, connection matrix.

2000 Mathematics Subject Classification Numbers: 33C99

1 Introduction

In many cases, hypergeometric functions have (or are defined by) Euler type integrals. In fact, the hypergeometric function associated to the configuration space of points on $\mathbb{P}^1$ has the following integral representation:

$$\int_{x_i}^{x_j} (t-x_1)^{\alpha_1}(t-x_2)^{\alpha_2}\cdots(t-x_n)^{\alpha_n}dt.$$

In this paper, instead of $\mathbb{P}^1$, we shall consider the configuration space of points on an elliptic curve.

Let C be an elliptic curve defined by $\mathbb{C}/\Gamma$, where $\Gamma := \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2$ is a lattice generated by ω_1 and ω_2. We shall denote a point on C by an equivalence class $\overline{x} \in C$ represented by $x \in \mathbb{C}$. For a triple $q = (\overline{x_0}, \overline{x_1}, \overline{x_2})$ of points on C, we denote by X_q by the punctured elliptic curve, that is, the open Riemann surface deprived of four points ($\overline{x_0}, \overline{x_1}, \overline{x_2}, \overline{0} \in C$). Our main objective is the integration of the multi-valued function g^{α}, where $\alpha \in \mathbb{C} \setminus \left(\frac{1}{2}\mathbb{Z} \cup \frac{1}{3}\mathbb{Z}\right)$ is a fixed complex number, whose branch points are $\overline{x_0}, \overline{x_1}, \overline{x_2}, \overline{0} \in C$. The function g is expressed by Weierstraß σ function:

$$g = \frac{\sigma(t-x_0)\sigma(t-x_1)\sigma(t-x_2)}{\sigma^3(t)\sigma(x_0)\sigma(x_1)\sigma(x_2)}.$$

It depends only on the equivalence class $\overline{t}$. So this function g is a meromorphic function on C holomorphic on X_q, which is a counterpart of rational functions on $\mathbb{P}^1$. Thus the function g^{α} can be seen as an analogue of the multi-valued function $t^{-3\alpha}(t-x_0)^{\alpha}(t-x_1)^{\alpha}(t-x_2)^{\alpha}$ on $\mathbb{P}^1$. However, it may depend on a representative x_0 (resp. x_1, x_2) of the equivalence class $\overline{x_0}$ (resp. $\overline{x_1}, \overline{x_2}$). To avoid this, we assume that $x_0 + x_1 + x_2 = 0$. The integration of $g^{\alpha}dt$ over a path $\Xi_{\mu,q}$ (mentioned below) lying on X_q

$$F_{\mu}(q) := \int_{\Xi_{\mu,q}} g^{\alpha}dt \tag{1}$$

is a multi-valued function with respect to q defined over the *configuration space* S (defined later), which is a subset of $\overline{S} := \{q = (\overline{x_0}, \overline{x_1}, \overline{x_2}) \mid x_0 + x_1 + x_2 = 0\} \subset C \times C \times C$.

This function F_{μ} has *singularities* : Put $D^{ij} := \left\{q = (\overline{x_0}, \overline{x_1}, \overline{x_2}) \in \overline{S} \mid x_i - x_j \in \Gamma\right\}$ and $D^{i}_{\infty} := \left\{q = (\overline{x_0}, \overline{x_1}, \overline{x_2}) \in \overline{S} \mid x_i \in \Gamma\right\}$; when q belongs to D^{ij}, the topological type of X_q differs from that of an punctured Riemann surface deprived of four points from C ; when q belongs to D^{i}_{∞}, the integrand g^{α} is divergent. Hence we define a domain S of the function F_{μ} by removing the singular loci D^{ij} and D^{i}_{∞} from $\overline{S}$.

We introduce an integration path $\Xi_{\mu,q}$ of (1), the so-called *twisted cycle*. Aomoto theory ([5]) tells us that a path of an Euler type integral can be regarded as a homology class with coefficients in the local system defined by the multi-valuedness of the integrand. In our case, the local system $\mathcal{L}_{X_q}$ is defined by the multi-valuedness of g^{α} and we denote by $\{\Xi_{\mu,q}\}_{\mu}$ generators of $H_1(X_q, \mathcal{L}_{X_q})$. In §3, we give the concrete description of $\{\Xi_{\mu,q}\}_{\mu}$ (Theorem 3.1). In §4, we calculate the intersection form among the twisted cycles (Theorem 4.1).

The function F_{μ} on S can be seen as a flat section of a certain vector bundle over S or, equivalently, a solution of a certain linear differential equation([8]) : The sheaf $\mathcal{S}$ of germs of functions defined by $\mathbb{C}$-linear combinations of $\{F_{\mu}\}_{\mu}$ is locally constant and $\mathcal{S} \otimes \mathcal{O}_S$ is a flat (trivial) vector bundle over S. The analytic continuation along a path γ in S induces the linear isomorphism between the stalks of $\mathcal{S}$ over the initial point and the terminal point of γ ; the corresponding matrix is said to be the *connection matrix*. In §5, we consider the paths in S connecting two points corresponding to a configuration $\left(\frac{\overline{\omega_i}}{2}, \frac{\overline{\omega_j}}{2}, \frac{\overline{\omega_k}}{2}\right)$ and another one given by interchanging two of the three points. For these paths, we calculate the connection matrices (Theorem 5.2).

Finally, we briefly mention the relation with the recent work of H.Watanabe([9][10]). He considers there a *fixed* configuration ; the two-torsion points are removed from elliptic curves, which are deformed by the moduli parameter. He focuses the action of modular transformations on such elliptic curves and studies that action on (co)homology groups of elliptic curves deprived of the two-torsion points with coefficients in certain local systems, which are of different kind of ours. In this context, he interprets the monodromy transformations of Gauß's hypergeometric function into the modular transformations.

Acknowledgments. The author would like to thank H.Watanabe and T.Mano for fruitful discussions. The author is sincerely grateful to K.Takamatsu for his guidance. The author would like to thank also Professors F.Kato and K.Fukaya for valuable comments and encouragements.

2 The local system

For a triple $q = (\overline{x_0}, \overline{x_1}, \overline{x_2})$ of points on $C = \mathbb{C}/\Gamma$, we denote by X_q an open Riemann surface deprived of four points $\overline{x_0}, \overline{x_1}, \overline{x_2}, \overline{0}$ from C. We assume that $x_0 + x_1 + x_2 = 0$ throughout this paper. We shall introduce a local system $\mathcal{L}_{X_q}$ which reflects the data of multi-valuedness of the function g^{α} defined as follows. Let g be the following holomorphic function on X_q:

$$g := \frac{\sigma(t - x_0)\sigma(t - x_1)\sigma(t - x_2)}{\sigma^3(t)\sigma(x_0)\sigma(x_1)\sigma(x_2)}.$$

Using the identity $\frac{\sigma'(t)}{\sigma(t)} = \zeta(t)$, we have $\frac{dg}{g} = (\zeta(t-x_0) + \zeta(t-x_1) + \zeta(t-x_2) - 3\zeta(t))dt$. We shall consider the following sheaf $\mathcal{L}_{X_q}$:

$$\mathcal{L}_{X_q} := \mathrm{Ker}\left(d - \alpha\frac{dg}{g} : \mathcal{O}_{X_q} \longrightarrow \Omega^1_{X_q}\right),$$

where $\alpha \in \mathbb{C} \setminus \left(\frac{1}{2}\mathbb{Z} \cup \frac{1}{3}\mathbb{Z}\right)$ is a fixed complex number. This sheaf has no global section on X_q. We take the open part $\overset{\circ}{X}_q$ in X_q such that $\mathcal{L}_{X_q}$ has global sections on $\overset{\circ}{X}_q$:

$$\overset{\circ}{X}_q := C \setminus \left(\overline{0x_0} \cup \overline{0x_1} \cup \overline{0x_2}\right),$$

where $\overline{0x_i}$ is the image of the segment $[0, x_i]$ in $\mathbb{C}$ through the natural projection $\mathbb{C} \longrightarrow C$.

Lemma 2.1. *1. The sheaf $\mathcal{L}_{X_q}$ is a local system of rank 1 over $\mathbb{C}$ whose local sections are generated by a branch of g^α.*

2. The sheaf $\mathcal{L}_{X_q}$ has a non-zero global section ς on $\overset{\circ}{X}_q$.

proof. 1. The function g^α is one of the solutions of the first-order ordinary linear differential equation $\left(d - \alpha\frac{dg}{g}\right)\varsigma = 0$. By the local theory of ordinary differential equations, local solutions of a first order linear differential equation are generated by a non-zero solution, whence the assertion.

2. We shall prove that a single-valued solution of the equation $\left(d - \alpha\frac{dg}{g}\right)\varsigma = 0$ on $\overset{\circ}{X}_q$ does exist. The logarithm of its solution is given by an integration of $\alpha\frac{dg}{g}$ over an arc. The single-valuedness is equivalent to the condition that the integration over any closed loop vanishes. Note that $\overset{\circ}{X}_q$ is homotopic to $S^1 \vee S^1$. In fact, we can pick two loops l_{ω_1}, l_{ω_2} on $\overset{\circ}{X}_q$ such that the inclusion map $l_{\omega_1} \cup l_{\omega_2} \hookrightarrow \overset{\circ}{X}_q$ is a deformation retract. That is the reason why we verify the integrations of $\frac{dg}{g}$ over those two loops l_{ω_1}, l_{ω_2} vanish. We denote by $I_i(x_0, x_1, x_2)$ these integrations:

$$I_i(x_0, x_1, x_2) := \int_{l_{\omega_i}} \frac{dg}{g} = \int_{l_{\omega_i}} (\zeta(t-x_0) + \zeta(t-x_1) + \zeta(t-x_2) - 3\zeta(t))dt.$$

Using the identity $\zeta'(t) = -\wp(t)$, we have $\frac{\partial I_i}{\partial x_j} = \int_{l_{\omega_i}} \wp(t - x_j)dt = \int_{l_{\omega_i}} \wp(t)dt$. This is a constant, which we denote by π_i. Then the equality $I_i(x_0, x_1, x_2) = \pi_i x_0 + \pi_i x_1 + \pi_i x_2 + I_i(0,0,0)$ holds. Note that $x_0 + x_1 + x_2 = 0$ (by the assumption) and $I_i(0,0,0) = 0$ (by its definition). Hence the integrations $I_i(x_0, x_1, x_2)$ vanish.

□

Now we pick a non-zero global section ς over $\overset{\circ}{X}_q$ and fix it once for all in the sequel. For a path $l : I \longrightarrow X_q$ with the initial point in $\overset{\circ}{X}_q$, we denote by ς_l the analytic continuation of ς along l.

Remark 2.2. *Note that X_q is homotopic to the bouquet $\underbrace{S^1 \vee \cdots \vee S^1}_{5\text{copies}}$. In fact, we can pick five loops l_0, l_1, l_2, l_{ω_1}, l_{ω_2} such that the inclusion map $l_0 \cup l_1 \cup l_2 \cup l_{\omega_1} \cup l_{\omega_2} \hookrightarrow X_q$ is a deformation retract. The loops l_{ω_1}, l_{ω_2} are the same ones as in the proof of lemma 2.1. The loop l_i is a closed curve around the point $\overline{x_i}$. The representation $\rho : \pi_1(X_q) \longrightarrow \mathbb{C}^*$ corresponding to the local system $\mathcal{L}_{X_q}$ is given by the formulae $\rho(l_i) = c$, $\rho(l_{\omega_i}) = 1$, where $c := e^{2\pi\sqrt{-1}\alpha}$.*

3 The twisted homology

We shall find generators of the *twisted homology group* $H_1(X_q, \mathcal{L}_{X_q})$.

The k-chain group $C_k(X_q, \mathcal{L}_{X_q})$ with coefficients in $\mathcal{L}_{X_q}$ is the complex vector space with basis $\sigma \otimes s_\sigma$, where $\sigma : \Delta \longrightarrow X_q$ is a singular k-simplex and s_σ is a section of $\mathcal{L}_{X_q}$ over σ. The first boundary operator $\partial : C_1(X_q, \mathcal{L}_{X_q}) \longrightarrow C_0(X_q, \mathcal{L}_{X_q})$ is given by $\partial(l \otimes s_l) = l(1) \otimes s_{l,l(1)} - l(0) \otimes s_{l,l(0)}$. The other boundary operators are given in similar fashions. The 1-st *twisted homology group* $H_1(X_q, \mathcal{L}_{X_q})$ is the 1-st homology of this complex, whose elements are called the *twisted cycles*. For two loops $l_{\omega_1}, l_{\omega_2}$ in the proof of lemma 2.1, we denote by $\Xi_{\omega_1,q}$ (resp. $\Xi_{\omega_2,q}$) the twisted cycle $l_{\omega_1} \otimes \varsigma_{l_{\omega_1}}$ (resp. $l_{\omega_2} \otimes \varsigma_{l_{\omega_2}}$), where ς is given in §2. We introduce other twisted cycles $\Xi_{(ij),q}$, which are the *regularizations* of the segments $\overline{x_i x_j}$:

$$\Xi_{(ij),q} := \frac{1}{c-1}\overline{S_\epsilon(x_i)} \otimes \varsigma_{\overline{S_\epsilon(x_i)}} + \overline{[x_i+\epsilon, x_j-\epsilon]} \otimes \varsigma_{\overline{[x_i+\epsilon, x_j-\epsilon]}} - \frac{1}{c-1}\overline{S_\epsilon(x_j)} \otimes \varsigma_{\overline{S_\epsilon(x_j)}},$$

where ϵ is a sufficiently small positive number; we denote by $[x_i+\epsilon, x_j-\epsilon]$ the oriented segment given by truncating each of both ends of the segment $[x_i, x_j]$ by the ϵ-circle centered at x_i or x_j; we denote by $S_\epsilon(x_i)$ a closed loop homotopic to the ϵ-circle centered at x_i both of whose initial and terminal points are at the initial point of the segment $[x_i+\epsilon, x_j+\epsilon]$, and $S_\epsilon(x_i)$ is defined in exactly the same way; we shall indicate by $^{-}$ an image through the natural projection $\mathbb{C} \longrightarrow C$. Note that if ϵ is sufficiently small, this definition is independent of ϵ.



Theorem 3.1 (generators of H_1).

1. *The chains* $\Xi_{(01),q}, \Xi_{(12),q}, \Xi_{(20),q}, \Xi_{\omega_1,q}, \Xi_{\omega_2,q}$ *are cycles and generate* $H_1(X_q, \mathcal{L}_{X_q})$.
2. *We have the relation* $\Xi_{(01),q} + \Xi_{(12),q} + \Xi_{(20),q} = 0$ *in* $H_1(X_q, \mathcal{L}_{X_q})$.

proof. The remark 2.2 tells us two facts. One is that the elements of $H_1(X_q, \mathcal{L}_{X_q})$ are represented by linear combinations of $l_0 \otimes \varsigma_{l_0}, l_1 \otimes \varsigma_{l_1}, l_2 \otimes \varsigma_{l_2}, l_{\omega_1} \otimes \varsigma_{l_{\omega_1}}, l_{\omega_2} \otimes \varsigma_{l_{\omega_2}}$. The other is that the topological euler number $\chi(X_q)$ equals 4. This implies that the dimension of $H_1(X_q, \mathcal{L}_{X_q})$ equals 4. We shall pick four linearly independent elements. The chain $\frac{1}{c-1} l_i \otimes \varsigma_{l_i} - \frac{1}{c-1} l_j \otimes \varsigma_{l_j}$ is a cycle homologous to $\Xi_{(ij),q}$. Immediately, we have the linear relation stated in this theorem. Then we obtain five elements of $H_1(X_q, \mathcal{L}_{X_q})$ subject to a single linear relation. The intersection form (*Theorem*4.1) in the next section tells us that any four of the five elements $\Xi_{(01),q}, \Xi_{(12),q}, \Xi_{(20),q}, \Xi_{\omega_1,q}, \Xi_{\omega_2,q}$ are linearly independent. □

4 The intersection form

The intersection form on twisted cycles is a bilinear form between $H_1\left(X_q, \mathcal{L}_{X_q}\right)$ and $H_1\left(X_q, \mathcal{L}^\vee_{X_q}\right)$, where $\mathcal{L}^\vee_{X_q}$ is the local system dual to $\mathcal{L}_{X_q}$; the local sections of $\mathcal{L}^\vee_{X_q}$ are generated by $g^{-\alpha}$ instead of g^α. To describe the intersection form concretely, we shall take the generators $\Xi^\vee_{(01),q}$, $\Xi^\vee_{(12),q}$, $\Xi^\vee_{(20),q}$, $\Xi^\vee_{\omega_1,q}$ and $\Xi^\vee_{\omega_2,q}$ of $H_1\left(X_q, \mathcal{L}^\vee_{X_q}\right)$ which are given by replacing ς (resp. c) by $\frac{1}{\varsigma}$ (resp. $\frac{1}{c}$) in the definitions of $\Xi_{(01),q}$, $\Xi_{(12),q}$, $\Xi_{(20),q}$, $\Xi_{\omega_1,q}$ and $\Xi_{\omega_2,q}$ respectively:

$$\varsigma \Longleftrightarrow \frac{1}{\varsigma}, \quad c \Longleftrightarrow \frac{1}{c} \quad ; \quad \Xi = \sum_\gamma a_\gamma(c)\gamma \otimes \varsigma_\gamma \Longleftrightarrow \Xi^\vee = \sum_\gamma a_\gamma\left(\frac{1}{c}\right)\gamma \otimes \frac{1}{\varsigma_\gamma},$$

where $\gamma \otimes \varsigma_\gamma$ is a twisted singular 1-simplex and $a_\gamma(X) \in \mathbb{C}(X)$.

We define the ordering on the index set $J := \{(01), (12), (20), \omega_1, \omega_2\}$ of the generators of H_1 :

$$(01) \prec (12) \prec (20) \prec \omega_1 \prec \omega_2.$$

Now we have

Theorem 4.1 (intersection form). *We assume* $\arg x_0 < \arg x_1 < \arg x_2$. *Then the intersection form between* $H_1\left(X_q, \mathcal{L}_{X_q}\right)$ *and* $H_1\left(X_q, \mathcal{L}^\vee_{X_q}\right)$ *is given by the matrix*

$$\left[\langle \Xi_{\mu,q}, \Xi^\vee_{\nu,q}\rangle\right]_{\mu,\nu\in J} = \begin{bmatrix} -\frac{c+1}{c-1} & \frac{1}{c-1} & \frac{c}{c-1} & 0 & 0 \\ \frac{c}{c-1} & -\frac{c+1}{c-1} & \frac{1}{c-1} & 0 & 0 \\ \frac{1}{c-1} & \frac{c}{c-1} & -\frac{c+1}{c-1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \end{bmatrix}.$$

In particular, the (2,2)-cofactor of this matrix is equal to $\frac{c^2+c+1}{(c-1)^2}$.

proof. The intersection number $\langle \Xi_{\mu,q}, \Xi^\vee_{\nu,q}\rangle$ can be calculated by summing up the local intersection numbers at each of the intersection points of the supports of $\Xi_{\mu,q}$ and of $\Xi^\vee_{\nu,q}$. The local intersection number $\langle \gamma_1 \otimes \varsigma_{\gamma_1}, \gamma_2 \otimes \frac{1}{\varsigma_{\gamma_2}}\rangle_p$ at the intersection point p of their supports is the product of the ordinary local intersection number $\langle \gamma_1, \gamma_2\rangle_p$ and $\frac{\varsigma_{\gamma_1}(p)}{\varsigma_{\gamma_2}(p)}$.

In the case of $\mu = \nu = (ij) \in \{(01), (12), (20)\}$:

We change the representative of the homology class $\Xi^\vee_{\nu,q}$ in such a way that its support meets that of $\Xi_{\mu,q}$ at two points p_i, p_j, which belong to the supports of $\overline{S_\epsilon(x_i)} \otimes \varsigma_{\overline{S_\epsilon(x_i)}}$ and of $\overline{S_\epsilon(x_j)} \otimes \varsigma_{\overline{S_\epsilon(x_j)}}$ respectively:

$$\begin{aligned} \langle \Xi_{\mu,q}, \Xi^\vee_{\nu,q}\rangle &= \frac{1}{c-1} \times (-1) \times \frac{\varsigma_{\overline{S_\epsilon(x_i)}}(p_i)}{\varsigma_{\overline{[x_i+\epsilon, x_j-\epsilon]}}(p_i)} - \frac{1}{c-1} \times 1 \times \frac{\varsigma_{\overline{S_\epsilon(x_j)}}(p_j)}{\varsigma_{\overline{[x_i+\epsilon, x_j-\epsilon]}}(p_j)} \\ &= \frac{1}{c-1} \times (-1) \times c - \frac{1}{c-1} \times 1 \times 1 = -\frac{c+1}{c-1}. \end{aligned}$$



In the case of $\mu \neq \nu$, $\mu, \nu \in \{(01), (12), (20)\}$:

We change the representative of the homology class $\Xi^{\vee}_{(jk),q}$ in such a way that its support meets that of $\Xi_{(ij),q}$ at a point p_j on the support of $\overline{S_\epsilon(x_j)} \otimes \varsigma_{\overline{S_\epsilon(x_j)}}$:

$$\langle \Xi_{(01),q}, \Xi^{\vee}_{(12),q} \rangle = \langle \Xi_{(12),q}, \Xi^{\vee}_{(20),q} \rangle = \langle \Xi_{(20),q}, \Xi^{\vee}_{(01),q} \rangle = -\tfrac{1}{c-1} \times (-1) \times \frac{\varsigma_{\overline{S_\epsilon(x_j)}}(p_j)}{\varsigma_{\overline{[x_i+\epsilon, x_j-\epsilon]}}(p_j)} = \frac{1}{c-1},$$

$$\langle \Xi_{(12),q}, \Xi^{\vee}_{(01),q} \rangle = \langle \Xi_{(20),q}, \Xi^{\vee}_{(12),q} \rangle = \langle \Xi_{(01),q}, \Xi^{\vee}_{(20),q} \rangle = \tfrac{1}{c-1} \times 1 \times \frac{\varsigma_{\overline{S_\epsilon(x_j)}}(p_j)}{\varsigma_{\overline{[x_i+\epsilon, x_j-\epsilon]}}(p_j)} = \frac{c}{c-1}.$$



In the case of $\mu, \nu \in \{\omega_1, \omega_2\}$:

The ordinary intersection form $\langle \bullet, \bullet \rangle : H_1(X_q, \mathbb{C}) \times H_1(X_q, \mathbb{C}) \longrightarrow \mathbb{C}$ is given by $\left[\langle l_{\omega_i}, l_{\omega_j} \rangle\right]_{i,j} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$, and $\frac{\varsigma_{l_{\omega_i}}(p)}{\varsigma_{l_{\omega_j}}(p)} = 1$ at the intersection point p of the supports of l_{ω_i} and of l_{ω_j}. Hence,

$$\left[\langle \Xi_{\omega_i,q}, \Xi^{\vee}_{\omega_j,q} \rangle\right]_{ij} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

In other cases:

We can change the representative of the homology class $\Xi^{\vee}_{\nu,q}$ in such a way that its support does not meet that of $\Xi_{\mu,q}$. □

5 The connection matrix

Put $\overline{S} := \left\{ q = (\overline{x_0}, \overline{x_1}, \overline{x_2}) \;\middle|\; x_0 + x_1 + x_2 = 0 \right\} \subset C \times C \times C$, $D^{ij} := \left\{ q \in \overline{S} \;\middle|\; x_i - x_j \in \Gamma \right\}$ and $D^i_\infty := \left\{ q \in \overline{S} \;\middle|\; x_i \in \Gamma \right\}$. We denote the *configuration space* by $S := \overline{S} \setminus \left(\bigcup_{i \neq j} D^{ij} \cup \bigcup_i D^i_\infty \right)$ and the point $\left(\frac{\overline{\omega_i}}{2}, \frac{\overline{\omega_j}}{2}, \frac{\overline{\omega_k}}{2} \right)$ on S by $q_{(ijk)}$, where $\omega_0 := -(\omega_1 + \omega_2)$. Let $\mathcal{S}$ be the sheaf over S of germs of functions defined by $\mathbb{C}$-linear combinations of $\left\{ F_\mu \right\}_\mu$; the stalk over $q_{(ijk)}$ is the vector space spanned by germs of $\left\{ \int_{\Xi_{\mu,q}} g^\alpha dt \right\}_{\mu \in \{(01),(20),\omega_1,\omega_2\}}$ at $q = q_{(ijk)}$.

In this chapter, we shall describe the linear isomorphisms among stalks of $\mathcal{S}$ over $q_{(012)}$, $q_{(210)}$ and $q_{(102)}$, of which the corresponding matrices are called the *connection matrices*, induced by the analytic continuations of $F_\mu(q) := \int_{\Xi_{\mu,q}} g^\alpha dt$ along four paths (given later) on S. These linear isomorphisms are identified with the isomorphisms among the homology groups $\left\{ H_1(X_q, \mathcal{L}_{X_q}) \right\}_{q \in \{q_{(012)}, q_{(210)}, q_{(102)}\}}$, which are induced by pulling back the fibration $\bigcup X_q \longrightarrow S$ along the paths on S. Put $\mathcal{H}(q) := H_1(X_q, \mathcal{L}_{X_q})$ for brevity.

We consider the paths in S, with $\overline{x_1}$ fixed, whose initial (resp. terminal) point is $q_{(012)}$ (resp. $q_{(210)}$). By their definitions, these paths are on the subset $\left\{q=(\overline{x_0},\overline{x_1},\overline{x_2})\in\overline{S}\;\middle|\;x_1=\frac{\omega_1}{2}\right\}$, which meets the singular loci at six points ; it meets D^{02} at four points $q^{1,0}_{(02)}, q^{-1,0}_{(02)}, q^{1,2}_{(02)}, q^{-1,2}_{(02)}$, meets both D^{01} and D^{2}_{∞} at the same point $q^{0,2}_{(02)}$, and meets both D^{12} and D^{0}_{∞} at the same point $q^{2,2}_{(02)}$, where

$$q^{m_1,m_2}_{(02)} := \left(\overline{\frac{\omega_0}{2}+\frac{m_1\omega_1+m_2\omega_2}{4}}, \overline{\frac{\omega_1}{2}}, \overline{\frac{\omega_2}{2}-\frac{m_1\omega_1+m_2\omega_2}{4}}\right).$$

We define four paths corresponding to the intersection points $q^{1,0}_{(02)}, q^{-1,0}_{(02)}, q^{1,2}_{(02)}, q^{-1,2}_{(02)}$ with D^{02} : The path $\gamma^{m_1,m_2}_{(02)}$ is defined by slightly deforming a path from $q_{(012)}$ to $q_{(210)}$ via $q^{m_1,m_2}_{(02)}$ in such a way that $\gamma^{m_1,m_2}_{(02)}$ avoids $q^{m_1,m_2}_{(02)}$:

$$\gamma^{m_1,m_2}_{(02)}(s) := \left(\overline{\frac{\omega_0}{2}+\frac{m_1\omega_1+m_2\omega_2}{2}s}, \overline{\frac{\omega_1}{2}}, \overline{\frac{\omega_2}{2}-\frac{m_1\omega_1+m_2\omega_2}{2}s}\right), \quad 0\le s\le\frac{1}{2}-\varepsilon,\ \frac{1}{2}+\varepsilon\le s\le 1,$$

where ε is a small positive number. In a similar fashion, we define the paths $\gamma^{0,1}_{(01)}, \gamma^{0,-1}_{(01)}, \gamma^{2,1}_{(01)}, \gamma^{2,-1}_{(01)}$, with $\overline{x_2}$ fixed, whose initial (resp. terminal) point is $q_{(012)}$ (resp. $q_{(102)}$) :

$$\gamma^{m_1,m_2}_{(01)}(s) := \left(\overline{\frac{\omega_0}{2}+\frac{m_1\omega_1+m_2\omega_2}{2}s}, \overline{\frac{\omega_1}{2}-\frac{m_1\omega_1+m_2\omega_2}{2}s}, \overline{\frac{\omega_2}{2}}\right), \quad 0\le s\le\frac{1}{2}-\varepsilon,\ \frac{1}{2}+\varepsilon\le s\le 1.$$

Note that the path $\gamma^{m_1,m_2}_{(ij)}$ can be realized as a composition of $\gamma^{1,0}_{(02)}$, $\gamma^{-1,0}_{(02)}$, $\gamma^{0,1}_{(01)}$ and $\gamma^{0,-1}_{(01)}$, so is a path, with $\overline{x_0}$ fixed, whose initial (resp. terminal) point is $q_{(012)}$ (resp. $q_{(021)}$). For example, $\gamma^{\pm1,2}_{(02)} = (\gamma^{0,1}_{(01)})^{\mp1}\circ(\gamma^{0,-1}_{(01)})^{\mp1}\circ(\gamma^{\pm1,0}_{(02)})^{\pm1}\circ(\gamma^{0,-1}_{(01)})^{\pm1}\circ(\gamma^{0,1}_{(01)})^{\pm1}$. (The following figure depicts the behavior of the representative $x^{m_1,m_2}_{0,(ij)}(s)$ of the $\overline{x_0}$-component of $\gamma^{m_1,m_2}_{(ij)}(s)$, where $q^{m_1,m_2}_{0,(ij)}$ denotes the representative of the $\overline{x_0}$-component of $q^{m_1,m_2}_{(ij)}$.)



For brevity, we denote by $\tau_{(ij)}$ the transposition (belonging to the symmetric group $\mathfrak{S}_3$) ; $\tau_{(ij)}(i)=j$, $\tau_{(ij)}(j)=i$ and $\tau_{(ij)}(k)=k$ for $k\neq i,j$.

We calculate the linear isomorphism $\left(\gamma^{m_1,m_2}_{(ij)}\right)_*$ between $\mathcal{H}\left(q_{(012)}\right)$ and $\mathcal{H}\left(q_{(\tau_{(ij)}(0),\tau_{(ij)}(1),\tau_{(ij)}(2))}\right)$ by the twisted Picard-Lefschetz formula ([7]) : The difference between a twisted cycle Ξ and the transformed one $\left(\gamma^{m_1,m_2}_{(ij)}\right)_*(\Xi)$ is equal to the product of the *vanishing cycle*, which vanishes at the singular point $q^{m_1,m_2}_{(ij)}$, and the constant factor which is calculated from the intersection numbers with the vanishing cycle and its *local monodromy* transformation.

Let $\Delta^{m_1,m_2}_{(ij)}(q)$ be the cycle vanishing at the singular point $q^{m_1,m_2}_{(ij)}$, as q tends to $q^{m_1,m_2}_{(ij)}$ along an extended path of $\gamma^{m_1,m_2}_{(ij)}$ that touches $q^{m_1,m_2}_{(ij)}$. We have the following :

Proposition 5.1 (vanishing cycles)**.**

$$\begin{aligned}
\Delta^{-1,0}_{(02)}(q_{(012)}) &= -\Xi_{(20),q_{(012)}} - \Xi_{\omega_1,q_{(012)}} - \Xi_{\omega_2,q_{(012)}},\\
\Delta^{1,0}_{(02)}(q_{(012)}) &= -\Xi_{(20),q_{(012)}} - \Xi_{\omega_2,q_{(012)}},\\
\Delta^{0,-1}_{(01)}(q_{(012)}) &= \Xi_{(01),q_{(012)}} - \Xi_{\omega_1,q_{(012)}} - \Xi_{\omega_2,q_{(012)}},\\
\Delta^{0,1}_{(01)}(q_{(012)}) &= \Xi_{(01),q_{(012)}} - \Xi_{\omega_1,q_{(012)}}.
\end{aligned}$$

proof. The vanishing cycle $\Delta^{m_1,m_2}_{(ij)}(q_{(012)})$ is a path on $X_{q_{(012)}}$ connecting two points $\overline{\frac{\omega_i}{2}}$, $\overline{\frac{\omega_j}{2}}$ depending on the path $\gamma^{m_1,m_2}_{(ij)}$. Replacing it by the linear combination of $\Xi_{(01),q_{(012)}}$, $\Xi_{(20),q_{(012)}}$, $\Xi_{\omega_1,q_{(012)}}$ and $\Xi_{\omega_2,q_{(012)}}$, we obtain the assertion. (See the following figures, in which the dotted lines indicate the vanishing cycles.)


$-\Xi_{\omega_1,q_{(012)}}$
$\frac{\omega_0}{2}$
$-\Xi_{\omega_2,q_{(012)}}$
$\Xi_{(20),q_{(012)}}$
$\Delta^{(-1,0)}_{(02)}(q_{(012)})$
ω_0
$\frac{\omega_0}{2}$
$\Delta^{1,0}_{(02)}(q_{(012)})$
$-\Xi_{\omega_2,q_{(012)}}$
$-\Xi_{(20),q_{(012)}}$
$-\omega_2$
$\frac{\omega_0}{2}$
$\Delta^{0,-1}_{(01)}(q_{(012)})$
ω_0
$-\Xi_{\omega_2,q_{(012)}}$
$\Xi_{(01),q_{(012)}}$
$-\Xi_{\omega_1,q_{(012)}}$
$-\omega_1$
$\Delta^{0,1}_{(01)}(q_{(012)})$
$\Xi_{(01),q_{(012)}}$
$-\Xi_{\omega_1,q_{(012)}}$
$\frac{\omega_0}{2}$

□

We have the common basis for $\mathcal{H}(q_{(012)})$ and $\mathcal{H}\left(q_{(\tau_{ij}(0),\tau_{ij}(1),\tau_{ij}(2))}\right)$: Put

$$\begin{aligned}
\Xi_{(kl)} &:= \frac{1}{c-1}\overline{S_\epsilon\left(\frac{\omega_k}{2}\right)} \otimes \varsigma_{\overline{S_\epsilon\left(\frac{\omega_k}{2}\right)}} + \overline{\left[\frac{\omega_k}{2}+\epsilon, \frac{\omega_l}{2}-\epsilon\right]} \otimes \varsigma_{\overline{\left[\frac{\omega_k}{2}+\epsilon, \frac{\omega_l}{2}-\epsilon\right]}} - \frac{1}{c-1}\overline{S_\epsilon\left(\frac{\omega_l}{2}\right)} \otimes \varsigma_{\overline{S_\epsilon\left(\frac{\omega_l}{2}\right)}}\\
\Xi_{\omega_k} &:= l_{\omega_k} \otimes \varsigma_{l_{\omega_k}},
\end{aligned}$$

and we define the ordering on the index set $J' := \{(01), (20), \omega_1, \omega_2\}$ of this basis :

$$(01) \prec (20) \prec \omega_1 \prec \omega_2.$$

Now we have the following :

Theorem 5.2 (connection matrices). *Let* $M(\gamma^{m_1,m_2}_{(ij)}) = \left[m(\gamma^{m_1,m_2}_{(ij)})_{\mu,\nu}\right]_{\mu,\nu\in J'}$ *be the* 4×4*-matrix representing the linear isomorphism* $\left(\gamma^{m_1,m_2}_{(ij)}\right)_*$*, that is,* $\left(\gamma^{m_1,m_2}_{(ij)}\right)_*(\Xi_\mu) = \sum_\nu m(\gamma^{m_1,m_2}_{(ij)})_{\mu,\nu}\Xi_\mu$*. Then,*

$$M(\gamma^{-1,0}_{(02)}) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ c & -c & c-1 & -c+1 \\ c & -c-1 & c & -c+1 \\ c & -c-1 & c-1 & -c+2 \end{bmatrix}, \qquad M(\gamma^{1,0}_{(02)}) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ c & -c & c-1 & 0 \\ 0 & 0 & 1 & 0 \\ c & -c-1 & c-1 & 1 \end{bmatrix},$$

$$M(\gamma^{0,-1}_{(01)}) = \begin{bmatrix} -c & 1 & -c+1 & c-1 \\ 0 & 1 & 0 & 0 \\ c+1 & -1 & c & -c+1 \\ c+1 & -1 & c-1 & -c+2 \end{bmatrix}, \qquad M(\gamma^{0,1}_{(01)}) = \begin{bmatrix} -c & 1 & 0 & c-1 \\ 0 & 1 & 0 & 0 \\ c+1 & -1 & 1 & -c+1 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

proof. Substituting the result of Proposition 5.1 for the twisted Picard-Lefschetz formula

$$\left(\gamma^{m_1,m_2}_{(ij)}\right)_*(\Xi) = \Xi + \frac{-c-1}{\left\langle \Delta^{m_1,m_2}_{(ij)}, \Delta^{m_1,m_2\vee}_{(ij)} \right\rangle} \left\langle \Xi, \Delta^{m_1,m_2\vee}_{(ij)} \right\rangle \Delta^{m_1,m_2}_{(ij)},$$

we obtain the assertion. □

Department of Mathematics
Faculty of Science
Kyoto University
Kitashirakawa-Oiwakechou,
Sakyouku, Kyoto 606-8502,
Japan
e-mail: koki@math.kyoto-u.ac.jp

References

[1] K.Aomoto, Equations aux différences linéaires et les intégrales des fonctions multiformes. I. Théorème d'existence, Proc. Japan Acad. 50(1974), 413–415.

[2] K.Aomoto, Équations aux différences linéaires et les intégrales des fonctions multiformes. II. Évanouissement des hypercohomologies et exemples, Proc. Japan Acad. 50(1974), 542–545.

[3] K.Aomoto, On vanishing of cohomology attached to certain many valued meromorphic functions, J. Math. Soc. Japan 27(1975), 248–255.

[4] K.Aomoto, Les équations aux différences linéaires et les intégrales des fonctions multiformes, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 22(1975), no. 3, 271–297.

[5] K.Aomoto, On the structure of integrals of power product of linear functions, Sci. Papers College Gen. Ed. Univ. Tokyo 27(1977), no. 2, 49–61

[6] K.Aomoto, M.Kita, Choukikakannsuuron (hypergeometric functions) (Japanese), Springer-Verlag Tokyo, 1994

[7] A.B.Givental, Twisted Picard Lefschetz formulas, translation in Funct. Anal. Appl. 22 (1988), no. 1, 10–18.

[8] K-K.Ito, T.Mano, The Elliptic Hypergeometric Functions Associated to the Configuration Space of Points on an Elliptic Curve II : Twisted de Rham cohomology, in preparation

[9] H.Watanabe, Transformation relations of matrix functions associated to the hypergeometric function of Gauss under modular transformations, J. Math. Soc. Japan 59 (2007), 113-126.

[10] H.Watanabe, Twisted homology and cohomology groups associated to the Wirtinger integral, J. Math. Soc. Japan 59 (2007), no. 4, 1067-1080.